AFRL-AFOSR-VA-TR-2017-0019



New Class of Excimer-Pumped Atomic Lasers (XPALS)

James Eden UNIVERSITY OF ILLINOIS CHAMPAIGN 506 S WRIGHT ST 364 HENRY ADMIN BLD URBANA, IL 61801-3620

01/27/2017 Final Report

DISTRIBUTION A: Distribution approved for public release.

Air Force Research Laboratory
AF Office Of Scientific Research (AFOSR)/RTB1

REPORT DOCUMENTATION PAGE

Form Approved OMS No. 0704-0188

The public reporting Ndenfor this COllection of information Is estinilled to avnge 1 hour per response, IndIldialgthe time for 1811ing InStructials, sewdlJng *15 LIIIg data sources, galhering and maintaining ...data needed. -9 and Al/lawing the collection of information indud019 suggestions for r.ducing the burden, to the Oepattment of e.e. eculive StMc:e Dilectorate (07()4.01118). Respondents Should be that stry OCher PIO" Islon of law, no person shall be subject to any penalty for Ialling to comply with ecolection of II> formation if iidoes not dis1liay *aitrently valid OMBoonIrd number.

PLEASE DONOT RETURN YOUR FORM TO THE ABOVE ORGANIZATION.				
1.REPORT DATE (DD-MM-YYYY) 12. REPORTTYPE 01/24/2017 Final	3. DATES COVERED (From - To) 06/01/2012 - 11/14/2015			
4. TITLE AND SUBTITLE	5a. CORALINUMBER			
New Class of Excimer-Pumped Atomic Lasers (XPALS)	FA9550-13-1-0006			
	5b.GRANT NUMBER			
	OB. GIVANT HOWIDER			
	Sc. f'ftOGRAM ELEMENT NUMBER			
6. AUTHOR(S)	Sd. PROJt:\. 1 NUMBER			
Eden, J.G., und Mironov, A. E.				
	5e. TASK NUMBER			
	JE. TASKNOWIDEK			
	5f. WORK UNTNUMBER			
7. PERFORMG ORGANIZATION NAME(S) AND ADDRESS(ES)	8. Pt:KFORMING ORGANIZATION			
Laboratory for Optical Physics and Engineering	REPORT NUMBER			
University or Illinois at Urbana-Champaign				
306 N. Wright SL				
Urbana, IL 61801-2918				
9. SPONSORING/MONTORING AGENCY NAME(S) AND ADDRESS(ES)	10. SPONSOR/MONITOR'S ACRONYM(S)			
USAF,AFRL DUNS 143574726				
AF OFFICE OF SCIENTIFIC RESEARCH				
875NORTH RANDOLPH STREET, RM 3112	11.5PONSOR/MOINOWS REPORT			
ARLINGTON VA 22203	NUMBER(S)			
SHANE C. LOMELIN 703-588-0196				
12. BTRIBUTION/AVAILABLITY STATEMENT				
DISTRIBUTION A: Distribution approved for public release.				
13. SUPPLEMENTARY NOTES				
13. SUPPLEMENTARY NOTES				
14. ABSTRACT				
This program has pursued the investigation or XPAL properties, and new pumping schemes, as well as modeling, and measuring critical				
photoionization and excited state-excited state reaction rates. We ore pleased to report that the main goal orthis program, the viability of nn atomic				
laser having a quantum efficiency greater thnn one, has been demonstrated. We believe this laser to represent a breakthrough in laser technology				
because the system appears to be scnlable in power but, more importBntly, the loser gain medium is cooled by the emission or every laser photon.				
The smallest energy derect that can exist between the pumped and lasing states in a three-level loser system has also been experimentally measured.				
The photoionization processes in nlkali-noble gas mixtures has been modeled and experimentally studied. Detailed modeling or XPAL and OPAL				
systems and plasma formation mechanisms during high power pumping has been performed.				
15.SUBJECTTERMS				
alkali lasers, optical pumping, laser medium cooling				
ABOTRACT	UMBER 19a. NAME OF RESPONSBLE PERSON			
a. REPORT b.ABSTRACT c.THS PAGE ABSTRACT OF	AGES CHIST COASCY			
	19b. TELEPHONE NUMBER (Includ11 area cod11)			
	217-244-7096			
	O: 000/D 0400			

FINAL REPORT OF AFOSR CONTRACT NO. FA9550-13-1-0006:

"New Class of Excimer-Pumped Atomic Lasers (XPALS)"

Prepared for
Susan Ecker
Contractor Support
Tech Marine Business (TMB)
Senior Program Analyst
Code 333
Office of Naval Research
Susan.Ecker.ctr@navy.mil

Prepared by
J. G. Eden and A. E. Mironov
Laboratory For Optical Physics and Engineering
Department of Electrical and Computer Engineering
University of Illinois
Urbana, IL 61801

November 2015

The focus of this 3 year AFOSR/HEL-JTO – sponsored research program was the investigation of XPAL properties, and new pumping schemes, as well as modeling, and measuring critical photoionization and excited state-excited state reaction rates in order to improve the performance of XPALs.

The first several experiments were conducted to measure the smallest energy defect that can exist between the pumped and lasing states in a three level laser system. This is a problem critical to the cooling of the laser medium by pumping it through the lower lying states. Results obtained during this program were described in detail in previous reports. In short, Rb-Xe gas mixtures were used to demonstrate the collapse of a three level laser when the thermal energy kT reaches approximately the energy defect between the laser and pumped levels.

Further experiments conducted at the University of Illinois have demonstrated the viability of an atomic laser having a quantum efficiency greater than one. We believe this laser to represent a breakthrough in laser technology because the system appears to be scalable in power but, more importantly, the laser gain medium is cooled by the emission of every laser photon.

Our experiments show that the suggested two color pumping scheme shown in fig. 1 allows the laser's efficiency to be increased by a factor of 1.7 (fig. 2). In addition, the




Figure 1. Schematic diagram of the laser system.




Figure 2. (Red circles) Dependence of D_2 line (780 nm) ASE on $E_1 + E_2$, the sum of the energies absorbed by the first and second pump pulses. For all measurements, [Rb]= $9.2*10^{14}$ cm⁻³, [Xe]= $1.8*10^{19}$ cm⁻³, λ_1 =759.95 nm, and λ_2 =794.76 nm. E_1 was fixed at 85 μ J and the time delay Δt was maintained at 8 ns; (Black Squares): Similar data were recorded when E_2 =0 and the laser is pumped only at 760 nm (Rb-Xe blue satellite).

pump absorption coefficient was increased by more than an order of magnitude, as compared to the classical XPAL pumping scheme. These results were published in Applied Physics Letters [1].

The experimental setup (Fig. 1) was described in detail in a previous report to the HEL-JTO. Briefly, a quartz cell filled with an alkali-noble gas mixture was pumped longitudinally with two dye lasers. The first pump pulse was used to prepare the system

and populated the upper laser level to a point slightly below the lasing threshold. The second (main) pump pulse created the population inversion and triggered lasing on the Rb D_2 line transition at 780 nm.

The most important feature of this pumping scheme is that the wavelength of the main pump was longer than the wavelength of the Rb D_2 line laser. In other words, the energy of the emitted photons was higher than the energy of the pump photons. This fact implies the extraction of thermal energy from the gain medium to compensate for the energy difference between the ASE and excitation photons.

It has been demonstrated (Fig. 2) that, in addition to the quantum efficiency being above one, the two-color pumped system exhibits higher slope efficiency as compared to the single color pumped case. It is important to note that the absolute efficiencies shown in Fig. 2 describe the ASE signals due to the specific experimental arrangement and, therefore, the actual D_2 line *laser* efficiency is significantly higher.

During the same reporting period, the kinetics of the higher energy excited states of Rb and Cs were studied in laser excitation experiments. In previous work, we have seen that the higher excited states are significantly populated by energy pooling processes when the D_1 or D_2 lines are strongly excited (e.g., optical pumping conditions). For the XPAL system, these higher excited states may be photoionized, resulting in loss of both input energy and metal atom number density. In a conventional DPAL system the higher excited states may react with hydrocarbons, producing metal hydride particles and carbon deposits.

Attempts were made to observe the ionization of Rb and Cs in the presence of 500 Torr of Ar. The D₂ lines were excited using pulsed laser intensities of approximately 0.5 MW/cm². A time-delayed, pulsed dye laser probe was used to detect metal vapor loss due to ionization. These measurements indicated that less than 1% of the neutral Rb or Cs was lost via ionization. Direct photoionization in the presence of 500 Torr of Ar, using 266 nm and 193 nm radation (P>1MW/cm²), was also found to be inefficient. Loss of ground state atoms was not detected, and there was negligible atomic line fluorescence.

Time-resolved fluorescence and laser pump-probe measurements were used to study the interactions of Rb(6p) with H_2 , CH_4 , and C_2H_6 . At room temperature, the total

removal, rate constants were found to be $k_{\rm H2}$ =(7.0±0.2)×10⁻¹⁰, $k_{\rm CH4}$ =(6.2±0.2)×10⁻¹⁰, and $k_{\rm C2H6}$ =(8.1±0.3)×10⁻¹⁰ cm³ s⁻¹. These values are consistent with earlier determinations. Electronic structure calculations were used to investigate the deactivation mechanisms. Quenching of Rb(6p) by H₂ proceeds via a curve crossing with the potential energy curve of the Rb(6s)+H₂ collision pair, while CH₄ and C₂H₆ quench via electronic-to-vibrational energy transfer. Measurements of ground state population recovery were used to estimate the fraction of the quenching that could be attributed to chemical reactions. For H₂ the reaction channel accounted for 12% of the total removal rate constant. Production of RbH was confirmed by observing the laser induced fluorescence (LIF) spectrum of the A¹ Σ^+ -X¹ Σ^+ transition. Reactive loss in the collisions with CH₄ and C₂H₆ accounted for 3% and 6% of the total removal rate constants. Surprisingly, searches for RbH produced by these reactions yielded negative results. Electronic structure calculations are currently being dedicated to examining the possible reactive channels for Rb(6p)+CH₄.

In summary, the main proposed milestones, and, in particular, the demonstration of the laser medium cooling with each laser photon produced, have all been achieved. The work conducted under this HEL-JTO program has demonstrated that the proposed two color pumping scheme of XPALs may be potentially used as the first high power laser system which is able to cool itself internally, thereby allowing its characteristics to be improved considerably as compared to other high power gas lasers.

We are grateful to the HEL-JTO for the support of this work.

List of publications:

- A. E. Mironov, W. Goldshlag and J. G. Eden, "Two color pumping of the Rb D₂ line laser (780 nm) through the photoassociation of Rb-Ar or Rb-Xe thermal pairs: Realization of a quantum efficiency above one", *Appl. Phys. Lett.*, 107, 041112 (2015).
- 2. J. D. Hewitt, T. J. Houlahan, Jr., J. E. Gallagher, D. L. Carroll, A. D. Palla, J. T. Verdeyen, G. P. Perram, and J. G. Eden, "Role of excited state photoionization in

- the 852.1 nm Cs laser pumped by Cs-Ar photoassociation," Appl. Phys. Lett., vol. 102, 111104 (March 18, 2013).
- 3. B. D. Barmashenko, S. Rosenwaks, and M. C. Heaven, Opt. Comm. 292, 123-125 (2013).
- 4. A. E. Mironov, W. Goldshlag, and J. G. Eden. "Alkali D₂ line laser optically pumped by two color free-free absorption," Proc. SPIE (Submitted)

AFOSR Deliverables Submission Survey

Response ID:7535 Data

1.

Report Type

Final Report

Primary Contact Email

Contact email if there is a problem with the report.

ccoxsey@illinois.edu

Primary Contact Phone Number

Contact phone number if there is a problem with the report

2172447096

Organization / Institution name

University of Illinois Urbana-Champaign

Grant/Contract Title

The full title of the funded effort.

NEW CLASS OF EXCIMER-PUMPED ATOMIC LASERS (XPALS)

Grant/Contract Number

AFOSR assigned control number. It must begin with "FA9550" or "F49620" or "FA2386".

FA9550-13-1-0006

Principal Investigator Name

The full name of the principal investigator on the grant or contract.

James Gary Eden

Program Officer

The AFOSR Program Officer currently assigned to the award

JASON A. MARSHALL, PhD

Reporting Period Start Date

06/01/2012

Reporting Period End Date

11/14/2015

Abstract

This program has pursued the investigation of XPAL properties, and new pumping schemes, as well as modeling, and measuring critical photoionization and excited state-excited state reaction rates. We are pleased to report that the main goal of this program, the viability of an atomic laser having a quantum efficiency greater than one, has been demonstrated. We believe this laser to represent a breakthrough in laser technology because the system appears to be scalable in power but, more importantly, the laser gain medium is cooled by the emission of every laser photon. The smallest energy defect that can exist between the pumped and lasing states in a three-level laser system has also been experimentally measured. The photoionization processes in alkali-noble gas mixtures has been modeled and experimentally studied. Detailed modeling of XPAL and DPAL systems and plasma formation mechanisms during high power pumping has been performed.

Distribution Statement

This is block 12 on the SF298 form.

Distribution A - Approved for Public Release

DISTRIBUTION A: Distribution approved for public release.

Explanation for Distribution Statement

If this is not approved for public release, please provide a short explanation. E.g., contains proprietary information.

SF298 Form

Please attach your SF298 form. A blank SF298 can be found here. Please do not password protect or secure the PDF The maximum file size for an SF298 is 50MB.

SF298.pdf

Upload the Report Document. File must be a PDF. Please do not password protect or secure the PDF. The maximum file size for the Report Document is 50MB.

HEL-JTO+REPORT+(AFOSR+0006)+2015+FINAL+REP.pdf

Upload a Report Document, if any. The maximum file size for the Report Document is 50MB.

Archival Publications (published) during reporting period:

- 1. A. E. Mironov, W. Goldshlag and J. G. Eden, "Two color pumping of the Rb D2 line laser (780 nm) through the photoassociation of Rb-Ar or Rb-Xe thermal pairs: Realization of a quantum efficiency above one", Appl. Phys. Lett., 107, 041112 (2015).
- 2. J. D. Hewitt, T. J. Houlahan, Jr., J. E. Gallagher, D. L. Carroll, A. D. Palla, J. T. Verdeyen, G. P. Perram, and J. G. Eden, "Role of excited state photoionization in the 852.1 nm Cs laser pumped by Cs-Ar photoassociation," Appl. Phys. Lett., vol. 102, 111104 (March 18, 2013).
- 3. B. D. Barmashenko, S. Rosenwaks, and M. C. Heaven, Opt. Comm. 292, 123-125 (2013).
- 4. A. E. Mironov, W. Goldshlag, and J. G. Eden. "Alkali D2 line laser optically pumped by two color free-free absorption," Proc. SPIE (Submitted)

New discoveries, inventions, or patent disclosures:

Do you have any discoveries, inventions, or patent disclosures to report for this period?

Νo

Please describe and include any notable dates

Do you plan to pursue a claim for personal or organizational intellectual property?

Changes in research objectives (if any):

None

Change in AFOSR Program Officer, if any:

None

Extensions granted or milestones slipped, if any:

None

AFOSR LRIR Number

LRIR Title

Reporting Period

Laboratory Task Manager

Program Officer

Research Objectives

Technical Summary

Funding Summary by Cost Category (by FY, \$K)

	Starting FY	FY+1	FY+2
Salary			
Equipment/Facilities			
Supplies			
Total			

Report Document

Report Document - Text Analysis

Report Document - Text Analysis

Appendix Documents

2. Thank You

E-mail user

Jan 24, 2017 15:32:58 Success: Email Sent to: ccoxsey@illinois.edu